

20dec01 07:41:54 User259284 Session D1538.2

SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1966-2002/JAN W2
 File 2:INSPEC 1969-2001/Dec W3
 (c) 2001 Institution of Electrical Engineers
 File 5:Biosis Previews(R) 1969-2001/Dec W3
 (c) 2001 BIOSIS
 File 6:NTIS 1964-2001/Dec W5
 (c) 2001 NTIS, Intl Cpyrght All Rights Res
 File 8:Ei Compendex(R) 1970-2001/Dec W3
 (c) 2001 Engineering Info. Inc.
 File 73:EMBASE 1974-2001/Dec W3
 (c) 2001 Elsevier Science B.V.
 File 987:TULSA (Petroleum Abs) 1965-2001/Dec W3
 (c) 2001 The University of Tulsa
 File 94:JICST-EPlus 1985-2001/Nov W2
 (c) 2001 Japan Science and Tech Corp(JST)
 File 35:Dissertation Abs Online 1861-2001/Dec
 (c) 2001 ProQuest Info&Learning
 File 144:Pascal 1973-2001/Dec W3
 (c) 2001 INIST/CNRS
 File 238:Abs. in New Tech & Eng. 1981-2001/Nov
 (c) 2001 Reed-Elsevier (UK) Ltd.
 File 105:AESIS 1851-2001/Jul
 (c) 2001 Australian Mineral Foundation Inc
 File 99:Wilson Appl. Sci & Tech Abs 1983-2001/Nov
 (c) 2001 The HW Wilson Co.
 File 58:GeoArchive 1974-2001/Nov
 (c) 2001 Geosystems
 File 34:SciSearch(R) Cited Ref Sci 1990-2001/Dec W4
 (c) 2001 Inst for Sci Info
 File 434:SciSearch(R) Cited Ref Sci 1974-1989/Dec
 (c) 1998 Inst for Sci Info
 File 292:GEOBASE(TM) 1980-2001/Dec
 (c) 2001 Elsevier Science Ltd.
 File 89:GeoRef 1785-2001/Dec B1
 (c) 2001 American Geological Institute
 File 65:Inside Conferences 1993-2001/Dec W3
 (c) 2001 BLDSC all rts. reserv.
 File 77:Conference Papers Index 1973-2001/Nov
 (c) 2001 Cambridge Sci Abs

S1 6358 FSE OR FAST() SPIN() ECHO OR FSEFT
 S2 1322 DRIVEN() EQUILIBRIUM OR DEFT
 S3 24 1AND2
 S4 9 RD S3 (unique items)
 S5 3 DRIVEN() EQUILIBRIUM() RADIOFREQUENC???/TI
 S6 6 SIMA/TI AND HADAMARD/TI
 S7 7 INTERACTIVE() FAST() SPIN() ECHO() IMAGING/TI
 S8 2 RD S7 (unique items)

4/9/9 (Item 1 from file: 77)
DIALOG(R)File 77:Conference Papers Index
(c) 2001 Cambridge Sci Abs. All rts. reserv.

4456633

Supplier Accession Number: 99-06018

V27N05

Cartilage imaging: Comparison of driven equilibrium Fourier transform
(DEFT) imaging with SPGR and FSE sequences

Lang, P.; Hargreaves, B.; Conolly, S.; Pauly, J.M.; Vandevenne, J.E.;
Lee, S.-U.; Gold, G.; Nishimura, D.

1999 European Congress of Radiology 9910013 Vienna (Austria) 7-12
Mar 1999

Agfa Gevaert N.V., Bracco s.p.a., GE Medical Systems - Europe, Mycomed
Imaging As, Philips Medical Systems, Schering AG, Siemens Medizinische
Technik

Springer Verlag, Tiergartenstr. 17, D-69121 Heidelberg, Germany; phone:
+49 6221 487 0; email: orders@springer.de; URL: www.springer.de, Abstracts
available. Contact Springer-Verlag for price. Paper No. 1429

Languages: ENGLISH

Descriptors: Multidisciplinary

Section Heading: MULTIDISCIPLINARY

Section Class Codes: 7000

4/9/4 (Item 4 from file: 155)
DIALOG(R) File 155:MEDLINE(R)

08372021 95241774 PMID: 7724816

Rapid MR imaging of the liver: comparison of twelve techniques for single breath-hold whole volume acquisition.

Naganawa S; Jenner G; Cooper TG; Potchen EJ; Ishigaki T

Department of Radiology, Michigan State University, USA.

Radiation medicine (JAPAN) Nov-Dec 1994, 12 (6) p255-61, ISSN 0288-2043 Journal Code: RAD

Languages: ENGLISH

Document type: Journal Article

Record type: Completed

Subfile: INDEX MEDICUS

Twelve magnetic resonance imaging pulse sequences for single breath-hold whole volume acquisition of the liver were evaluated on volunteers. Liver and spleen contrast to noise ratio (C/N), overall image quality, and grade of artifacts were compared. The 12 sequences included T2-weighted **fast spin echo (FSE)** with or without fat suppression (FS), fast multiplanar spoiled gradient recalled imaging (FMPSPGR), fast gradient recalled imaging without preparation pulses (FGR), FGR with inversion recovery preparation pulse nulling the liver or fat (IR-FGR-L and IR-FGR-F), FGR with **driven equilibrium** preparation pulse (DE-FGR), single shot moderately or heavily T2-weighted spin echo echo planar imaging (SE-EPI-mT2 and SE-EPI-hT2), multi-shot moderately T2-weighted spin echo echo planar imaging (multi-shot SE-EPI-mT2), inversion recovery EPI, and gradient echo EPI. In the quantitative analysis, **FSE + FS** showed a significantly higher C/N ratio than the others ($p < 0.05$). In the qualitative evaluation, DE-prepFGR, and single and multi-shot SE-EPI-mT2 had good results, as did **FSE** and **FSE + FS**. Further studies should be conducted to determine whether or not these breath-hold sequences can obviate current conventional non-breath-hold sequences.

Tags: Comparative Study; Human

4/9/7 (Item 1 from file: 34)
DIALOG(R)File 34:SciSearch(R) Cited Ref Sci
(c) 2001 Inst for Sci Info. All rts. reserv.

04250788 Genuine Article#: RR687 Number of References: 11
Title: HIGH-CONTRAST AND FAST 3-DIMENSIONAL MAGNETIC-RESONANCE-IMAGING AT
HIGH FIELDS

Author(s): LEE JH; GARWOOD M; MENON R; ADRIANY G; ANDERSEN P; TRUWIT CL;
UGURBIL K

Corporate Source: UNIV MINNESOTA,CTR MAGNET RESONANCE RES,SCH MED,385 E
RIVER RD/MINNEAPOLIS//MN/55455; UNIV MINNESOTA,CTR MAGNET RESONANCE
RES,SCH MED/MINNEAPOLIS//MN/55455; UNIV MINNESOTA,DEPT CHEM,DEPT
RADIOL/MINNEAPOLIS//MN/55455

Journal: MAGNETIC RESONANCE IN MEDICINE, 1995, V34, N3 (SEP), P308-312
ISSN: 0740-3194

Language: ENGLISH Document Type: NOTE

Geographic Location: USA

Subfile: SciSearch; CC CLIN--Current Contents, Clinical Medicine

Journal Subject Category: RADIOLOGY & NUCLEAR MEDICINE

Abstract: A new three-dimensional imaging strategy based on magnetization prepared ultrafast gradient recalled echo technique that demonstrates pronounced T-1 contrast at high fields is introduced, High-resolution three-dimensional image sets of human brain showing high contrast between white and gray matter areas are presented, The ratio of contrast-to-noise was examined as a function of the relevant parameters in the imaging sequence; calculations based on high-field T-1 values as well as the experimental data demonstrated that maximal contrast-to-noise ratio is attained under the same magnetization preparation conditions both for cortical and subcortical gray matter relative to white matter, leading to approximately equivalent appearance of all gray matter areas in the same image. In addition, the images displayed clear visualization of subtle anatomical structures such as the subthalamic nuclei (ventral tier nuclei, dorsomedial nucleus, and pulvinar) and mammillothalamic tracts.

Descriptors--Author Keywords: MRI ; PULSE SEQUENCES ; MODIFIED **DRIVEN**
EQUILIBRIUM FOURIER TRANSFORM ; HIGH T-1 CONTRAST